



# A global uniqueness result for acoustic tomography of moving fluid

Alexey Agaltsov

## ► To cite this version:

Alexey Agaltsov. A global uniqueness result for acoustic tomography of moving fluid. 2015. hal-01122389

**HAL Id: hal-01122389**

**<https://hal.science/hal-01122389>**

Preprint submitted on 3 Mar 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A global uniqueness result for acoustic tomography of moving fluid

*A. D. Agaltsov*<sup>1,2</sup>

We consider a model time-harmonic wave equation of acoustic tomography of moving fluid in an open bounded domain in dimension  $d \geq 2$ . We give global uniqueness theorems for related inverse boundary value problem at fixed frequency.

**Keywords:** inverse boundary value problems, time-harmonic wave equation, acoustic tomography

**Subjects:** partial differential equations, mathematical physics

**AMS classification:** 35R30 (Inverse problems), 35Q35 (PDEs in connection with fluid mechanics)

## 1 Introduction

Consider the operator

$$L_{A,V} = -\Delta - 2iA(x) \cdot \nabla + V(x), \quad (1)$$

where  $\Delta$  is the standard Laplacian,  $x \in D$ ,  $A \in W^{1,\infty}(D, \mathbb{R}^d)$ ,  $V \in L^\infty(D, \mathbb{R})$ ,  $D$  is an open bounded domain in  $\mathbb{R}^d$  ( $d \geq 2$ ). In the present article we study an inverse boundary value problem for the equation  $L_{A,V}\psi = 0$  in  $D$ .

As in [AN], [RW], [RE] we consider the equation  $L_{A,V}\psi = 0$  as a model equation for a time-harmonic ( $e^{-i\omega t}$ ) pressure  $\psi$  in moving fluid. In this setting

$$A(x) = \frac{\omega}{c^2(x)}v(x), \quad V(x) = -\frac{\omega^2}{c^2(x)},$$

where  $v$  is the fluid velocity vector,  $c$  is the sound speed,  $\omega$  is the frequency.

Suppose that 0 is not a Dirichlet eigenvalue for operator  $L_{A,V}$  in  $D$ . Then the Dirichlet problem

$$\begin{cases} L_{A,V}\psi = 0 & \text{in } D, \\ \psi|_{\partial D} = f, \end{cases} \quad (2)$$

---

<sup>1</sup>Centre de Mathématiques Appliquées, Ecole Polytechnique  
Route de Saclay

91128 PALAISEAU Cedex, France

<sup>2</sup>Lomonosov Moscow State University,

GSP-1, Leninskie Gory

119991 MOSCOW, Russia

email: agalets@gmail.com

is uniquely solvable for  $\psi \in H^1(D)$  for any  $f \in H^{1/2}(\partial D)$ . The Dirichlet-to-Neumann map  $\Lambda_{A,V}$  sends  $f \in H^{1/2}(\partial D)$  to  $\Lambda_{A,V}f \in H^{-1/2}(\partial D)$  defined by the formula

$$\Lambda_{A,V}f = \frac{\partial \psi}{\partial \nu} \Big|_{\partial D} + i(A \cdot \nu)f, \quad (3)$$

where  $\nu$  is the unit exterior normal to  $\partial D$  and  $\frac{\partial \psi}{\partial \nu} \Big|_{\partial D} \in H^{-1/2}(\partial D)$  can be defined, in particular, by the following formula:

$$\langle \frac{\partial \psi}{\partial \nu} \Big|_{\partial D}, u \rangle = \int_D \left( \nabla \psi(x) \nabla \tilde{u}(x) - 2i\tilde{u}(x)A(x) \nabla \psi(x) + \tilde{u}(x)V(x)\psi(x) \right) dx, \quad (4)$$

for  $u \in H^{1/2}(\partial D)$  and arbitrary  $\tilde{u} \in H^1(D)$  with  $\tilde{u}|_{\partial D} = u$ . Note that since  $\psi$  satisfies  $L_{A,V}\psi = 0$ , the right hand side of the above formula doesn't depend on the choice of  $\tilde{u}$ .

The inverse boundary value problem for equation  $L_{A,V}\psi = 0$  in  $D$  consists in finding  $A, V$  from  $\Lambda_{A,V}$ . In the case when coefficients  $A, V$  can be complex-valued there is an obstruction to the unique solvability of this problem caused by the gauge invariance of the map  $\Lambda_{A,V}$  with respect to the gauge transformations

$$\begin{cases} A \rightarrow A + \nabla \varphi, \\ V \rightarrow V - i\Delta \varphi + (\nabla \varphi)^2 + 2A \nabla \varphi, \end{cases}$$

where  $\varphi \in W^{2,\infty}(D, \mathbb{C})$ ,  $\varphi|_{\partial D} = 0$ , see, e.g., [KU] ( $d \geq 3$ ), [GT] ( $d = 2$ ).

However, in the case of real-valued coefficients  $A, V$  there is no gauge non-uniqueness as it was observed, for example, in [AN].

In addition, in general case, under some regularity assumptions on  $\partial D, A$  and  $V$ , the Dirichlet-to-Neumann map  $\Lambda_{A,V}$  uniquely determines the two-form  $dA$  and the function  $q$  in  $D$  and the tangential component of  $A$  on  $\partial D$ , where

$$\begin{aligned} dA &= \sum_{1 \leq k < l \leq d} \left( \frac{\partial A^l}{\partial x_k} - \frac{\partial A^k}{\partial x_l} \right) dx_k \wedge dx_l, \\ q &= V + i \nabla \cdot A - A \cdot A, \end{aligned} \quad (5)$$

and  $A = (A^1, \dots, A^d)$ .

In particular, it was shown in [KU] that in dimension  $d \geq 3$  the map  $\Lambda_{A,V}$  uniquely determines  $dA$  and  $q$  in  $D$  if  $A \in L^\infty(D, \mathbb{C}^d)$  and  $V \in L^\infty(D, \mathbb{C})$ . And in dimension  $d = 2$  it was shown in [GT] that if  $D$  is a smooth Riemann surface with boundary (in particular, if  $D$  is a planar domain with  $\partial D \in C^\infty$ ) then the map  $\Lambda_{A,V}$  uniquely determines  $dA$  and  $q$  provided that  $A \in W^{2,p}(D, \mathbb{R}^d)$ ,  $V \in W^{1,p}(D, \mathbb{C})$ ,  $p > 2$ .

In addition, concerning the identifiability of tangential components of  $A$  on the boundary, it was proved in [BS] that if  $\partial D \in C^1$  ( $d \geq 3$ ) or  $\partial D \in C^{1,\alpha}$ ,  $\alpha \in (0, 1)$  ( $d = 2$ ) and if  $A \in C(\overline{D}, \mathbb{C}^d)$ ,  $V \in L^\infty(D, \mathbb{C})$  then  $\Lambda_{A,V}$  uniquely determines  $A - \nu(A \cdot \nu)$  on  $\partial D$ , where  $\nu$  is the unit exterior normal field to  $\partial D$ .

In the present article we combine the aforementioned results in order to obtain the following global uniqueness results in the case when coefficients  $A, V$  are real-valued.

**Theorem 1.** *Let  $D$  be a bounded simply connected domain with path connected boundary in  $\mathbb{R}^d$  ( $d \geq 3$ ) with  $\partial D \in C^1$ . Let  $A_1, A_2 \in W^{1,\infty}(D, \mathbb{R}^d)$  and  $V_1, V_2 \in L^\infty(D, \mathbb{R})$ . If  $\Lambda_{A_1, V_1} = \Lambda_{A_2, V_2}$ , then  $A_1 = A_2, V_1 = V_2$ .*

**Theorem 2.** *Let  $D$  be a bounded simply connected domain in  $\mathbb{R}^2$  with  $\partial D \in C^\infty$ . Let  $A_1, A_2 \in W^{2,p}(D, \mathbb{R}^d)$  and  $V_1, V_2 \in W^{1,p}(D, \mathbb{R})$  with  $p > 2$ . If  $\Lambda_{A_1, V_1} = \Lambda_{A_2, V_2}$  then  $A_1 = A_2$  and  $V_1 = V_2$ .*

Theorems 1 and 2 are proved in Section 3. In Section 2 we present formulas and equations for finding  $A, V$  from  $dA, q, A - \nu(A \cdot \nu)|_{\partial D}$ .

## 2 Formulas and equations for finding $A, V$

In this section we suppose that  $D$  is a bounded contractible domain with path connected  $C^2$  boundary in  $\mathbb{R}^d$  ( $d \geq 2$ ). By contractibility we mean that there exists  $F \in C^2(D \times [0, 1], D)$  such that  $F_0 \equiv \tilde{x}, F_1 = \text{id}_D$ , where  $F_t(x) = F(x, t)$ ,  $\tilde{x}$  is some fixed point in  $D$  and  $\text{id}_D$  is the identity mapping on  $D$ . We also suppose that  $A \in W^{2,\infty}(D, \mathbb{R}^d), V \in L^\infty(D, \mathbb{R})$ . Given  $dA, q$  as in (5) in  $D$  and  $A - \nu(A \cdot \nu)$  on  $\partial D$ , we can find  $A, V$  in the following way:

1. Define  $\tilde{A} = (\tilde{A}^1, \dots, \tilde{A}^d) \in W^{1,\infty}(D, \mathbb{R}^d)$  by the formula

$$\tilde{A}^k = \sum_{i < j} \int_0^1 \left( \frac{\partial F_t^i}{\partial t} \frac{\partial F_t^j}{\partial x_k} - \frac{\partial F_t^j}{\partial t} \frac{\partial F_t^i}{\partial x_k} \right) \left( \frac{\partial A^j}{\partial x_i} \circ F_t - \frac{\partial A^i}{\partial x_j} \circ F_t \right) dt, \quad (6)$$

where  $k = 1, \dots, d; A = (A^1, \dots, A^d), F_t = (F_t^1, \dots, F_t^d)$  and  $\circ$  denotes the composition of maps, i.e.  $\frac{\partial A^j}{\partial x_i} \circ F_t(y) = \frac{\partial A^j}{\partial x_i}(F_t(y)), y \in D$ .

2. Fix  $x^0 \in \partial D$ . Define  $\varphi_0 \in C^1(\partial D)$  by the formula

$$\varphi_0(x) = \sum_{k=1}^d \int_{x^0}^x (A_\tau^k(y) - \tilde{A}_\tau^k(y)) dy_k, \quad x \in \partial D, \quad (7)$$

where  $A_\tau = A - \nu(A \cdot \nu), \tilde{A}_\tau = \tilde{A} - \nu(\tilde{A} \cdot \nu), A_\tau = (A_\tau^1, \dots, A_\tau^d), \tilde{A}_\tau = (\tilde{A}_\tau^1, \dots, \tilde{A}_\tau^d), \nu$  is the unit exterior normal field to  $\partial D$  and integration is over an arbitrary  $C^1$  curve on  $\partial D$  linking  $x^0$  to  $x$ .

3. Find the unique generalized solution  $\varphi \in W^{2,\infty}(D, \mathbb{R})$  to

$$\begin{cases} \Delta \varphi = \text{Im } q - \nabla \cdot \tilde{A} & \text{in } D, \\ \varphi|_{\partial D} = \varphi_0. \end{cases} \quad (8)$$

4. Coefficients  $A, V$  are given by the following formulas:

$$\begin{cases} A = \tilde{A} + \nabla \varphi, \\ V = q - i \Delta \varphi - i \nabla \cdot \tilde{A} + \tilde{A} \cdot \tilde{A} + 2 \tilde{A} \cdot \nabla \varphi + (\nabla \varphi)^2. \end{cases}$$

This algorithm will be justified in Section 4.

### 3 Proofs of Theorems 1, 2

We will prove Theorems 1 and 2 simultaneously. Let  $D$ ,  $A_1$ ,  $A_2$ ,  $V_1$ ,  $V_2$  satisfy the conditions of Theorem 1 (resp. Theorem 2) and suppose that  $\Lambda_{A_1, V_1} = \Lambda_{A_2, V_2}$ .

Using Theorem 1.1 of [BS] we obtain that

$$(A_1 - \nu(A_1 \cdot \nu))|_{\partial D} = (A_2 - \nu(A_2 \cdot \nu))|_{\partial D}, \quad (9)$$

where  $\nu$  is the unit exterior normal field to  $\partial D$ . Using Theorem 1.1 of [KU] (resp. Theorem 1.1 of [GT]) we get

$$dA_1 = dA_2 \text{ in } D, \quad (10)$$

$$q_1 = q_2 \text{ in } D, \quad (11)$$

where

$$dA_j = \sum_{1 \leq k < l \leq d} \left( \frac{\partial A_j^l}{\partial x_k} - \frac{\partial A_j^k}{\partial x_l} \right) dx_k \wedge dx_l, \\ q_j = V_j + i \nabla \cdot A_j - A_j \cdot A_j,$$

where  $A_j = (A_j^1, \dots, A_j^d)$ ,  $j = 1, 2$ .

Since the domain  $D$  is simply connected it follows from (10) that there exists  $\varphi \in W^{2,\infty}(D, \mathbb{R})$  such that

$$A_1 - A_2 = \nabla \varphi \text{ in } D. \quad (12)$$

In dimension  $d = 2$  it follows from simple connectedness of  $D$  and from smoothness of  $\partial D$  that  $\partial D$  is path connected. Formulas (9), (12) and path connectedness of  $\partial D$  imply that  $\varphi$  is constant on  $\partial D$ .

Using (11), (12) we obtain that

$$V_1 - V_2 = -i\Delta\varphi - (\nabla\varphi)^2 + 2A_1\nabla\varphi \text{ in } D.$$

Taking the imaginary part of this equation we obtain the equation  $\Delta\varphi = 0$  in  $D$ . Since  $\varphi$  is constant on  $\partial D$  and  $\varphi \in W^{2,\infty}(D)$  it follows that  $\varphi$  is constant in  $\overline{D}$ . Hence  $A_1 = A_2$  and  $V_1 = V_2$ . Theorems 1 and 2 are proved.

### 4 Justification of the algorithm of Section 2

It follows from formula (6) that  $d\tilde{A} = dA$ . More precisely, if we denote by  $F^*dA$  the pullback of the form  $dA$  by the map  $F$  and by  $\iota_{\partial_t}$  we denote the interior product with the vector field  $\frac{\partial}{\partial t}$  on  $\{(x, t) \in D \times [0, 1]\}$ , then

$$\sum_{k=1}^d \tilde{A}^k dx_k = \int_0^1 (\iota_{\partial_t} F^* dA) dt,$$

and the equality  $d\tilde{A} = dA$  follows from the Cartan magic formula  $\mathcal{L}_{\partial_t} = d \circ \iota_{\partial_t} + \iota_{\partial_t} \circ d$ , where  $\mathcal{L}_{\partial_t}$  is the Lie derivative along  $\frac{\partial}{\partial t}$ ,  $d$  is the exterior derivative on  $\{(x, t) \in D \times [0, 1]\}$  and  $\circ$  denotes the composition of maps.

Hence we can define  $\varphi \in W^{2,\infty}(D, \mathbb{R})$  by the formula

$$\varphi(x) = \int_{\tilde{x}}^x \sum_{k=1}^d (A_k - \tilde{A}_k) dx_k, \quad x \in \overline{D},$$

where  $\tilde{x} \in D$  is some fixed point and integration is over an arbitrary  $C^1$  curve in  $D$  linking  $\tilde{x}$  to  $x$ . Then  $\nabla \varphi = A - \tilde{A}$  in  $\overline{D}$  and this implies that  $\varphi|_{\partial D}$  differs by constant from  $\varphi_0$  defined in (7). We also obtain from (5) the equation

$$V = q - i\Delta\varphi - i\nabla \cdot \tilde{A} + \tilde{A} \cdot \tilde{A} + 2\tilde{A} \cdot \nabla\varphi + (\nabla\varphi)^2 \quad \text{in } D.$$

Taking into account that  $V$  is real-valued and separating the imaginary part in the latter equation we obtain (8). Since  $\text{Im } q$  and  $\nabla \cdot \tilde{A}$  belong to  $L^\infty(D, \mathbb{R})$ , the problem (8) is uniquely solvable for  $\varphi \in W^{2,\infty}(D, \mathbb{R})$ . Thus, the algorithm of Section 2 is justified.

## 5 Acknowledgements

This work was fulfilled within the framework of research carried out under the supervision of Prof. R.G. Novikov.

## 6 References

- [AN] A.D. Agaltsov, R.G. Novikov, Riemann-Hilbert problem approach for two-dimensional flow inverse scattering, *J. Math. Phys.* **55** (10), 2014, 103502
- [BS] R. M. Brown, M. Salo, Identifiability at the boundary for first-order terms, *Appl. Anal.* **85** (6–7), 2006, 735–749
- [GT] C. Guillarmou, L. Tzou, Identification of a connection from Cauchy data on a Riemann surface with boundary, *Geom. Funct. Anal.* **21** (2), 2011, 393–418
- [KU] K. Krupchyk, G. Uhlmann, Uniqueness in an inverse boundary problem for a magnetic Schrödinger operator with a bounded magnetic potential, *Comm. Math. Phys.* **327** (3), 2014, 993–1009
- [RW] D. Roussef, K. B. Winters, Two-dimensional vector flow inversion by diffraction tomography, *Inverse Problems* **10**, 1994, 687–697
- [RE] M. N. Rychagov, H. Ermert, Reconstruction of fluid motion in acoustic diffraction tomography, *J. Acoust. Soc. Am.* **99** (5), 1996, 3029–3035